

basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

NATIONAL SENIOR CERTIFICATE/ *NASIONALE SENIOR SERTIFIKAAT*

GRADE/GRAAD 11

MATHEMATICS P1/WISKUNDE V1

NOVEMBER 2016

MEMORANDUM

MARKS/PUNTE: 150

**This memorandum consists of 18 pages.
Hierdie memorandum bestaan uit 18 bladsye.**

**DEPARTMENT OF BASIC
EDUCATION
PRIVATE BAG X800, PRETORIA 0001**

2016 -11- 11

**APPROVED MARKING GUIDELINE
PUBLIC EXAMINATION**

*D.M.
12/11/2016*

NOTE:


- If a candidate answered a question TWICE, mark only the FIRST attempt.
- If a candidate crossed out an answer and did not redo it, mark the crossed-out answer.
- Consistent accuracy applies to ALL aspects of the marking memorandum(If a learner makes a mistake, the mistake has to be followed up. Stop marking the question if the learner commits the second mistake)
- It is unacceptable to assume values/answers in order to solve a problem.
- Penalise once for incorrect rounding off.




LET WEL:


- As 'n kandidaat 'n vraag TWEE keer beantwoord het, sien slegs die EERSTE poging na.
- As 'n kandidaat 'n antwoord deurgehaal en nie oorgedoen het nie, sien die deurgehaalde antwoord na.
- Volgehoue akkuraatheid is op ALLE aspekte van die memorandum van toepassing(as 'n leerder 'n fout gemaak het, moet die fout opgevolg word. Hou op om die vraag na te sien as die leerder 'n tweede fout maak)
- Dit is onaanvaarbaar om waardes/antwoorde te veronderstel om 'n probleem op te los.
- Penaliseer een keer vir inkorrekte afronding

QUESTION/VRAAG 1

1.1.1	$3x^2 - 5x - 1 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-(-5) \pm \sqrt{(-5)^2 - 4(3)(-1)}}{2(3)}$ $= \frac{5 \pm \sqrt{37}}{6}$ $x = 1,85 \text{ or } x = -0,18$ <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> Incorrect rounding off, max $\frac{2}{3}$ marks </div>	✓ substitution in the correct quadratic formula /vervanging in die korrekte formule ✓ answer/antwoord ✓ answer/antwoord (3)
1.1.2	$x^2 - 6x + 8 = 0$ $(x - 4)(x - 2) = 0$ $x = 4 \text{ or } x = 2$	✓ factors/faktore OR Substitution in the correct quadratic formula/ vervanging in die korrekte kwadratiese formule ✓ $x = 4$ ✓ $x = 2$ (3)




<p>1.1.3 Option/Opsie 1</p> $4x - 2x^2 < 0$ $2x(2-x) < 0$ $x < 0 \text{ or } x > 2$ 	OR/OF	<p>Option/Opsie 2</p> $4x - 2x^2 < 0$ $-2x^2 + 4x < 0$ $2x^2 - 4x > 0$ $x(2x - 4) > 0$ $x < 0 \text{ or } x > 2$  OR 	<p>✓ factors/faktore ✓ method/metode ✓✓ $x < 0$ or $x > 2$</p> <p>Maximum $\frac{3}{4}$ for incorrect notation (4)</p> <p>Maksimum $\frac{3}{4}$ vir verkeerde notasie</p>
---	--------------	---	--



D-m


1.1.4	$2^{3x+1} + 2^{3x} = 12$ $2^{3x} [2^1 + 1] = 12$ $2^{3x} \cdot 3 = 12$ $2^{3x} = 4$ $2^{3x} = 2^2$ $3x = 2$ $\therefore x = \frac{2}{3}$	✓ common/gemene factor ✓ simplification/vereenv. ✓ equating/gelykst exponents ✓ answer/antw. (4)
1.1.5	$\sqrt{x-1} + 3 = x - 4$ $\sqrt{x-1} = x - 4 - 3$ $x-1 = (x-7)^2$ $x-1 = x^2 - 14x + 49$ $x^2 - 15x + 50 = 0$ $(x-5)(x-10) = 0$ $x \neq 5 \text{ or } x = 10$	✓ isolate/soleer ✓ sign/teken ✓ squaring/kwadr both sides ✓ std vorm/stand vorm ✓ factors/fakt ✓ $x \neq 5$ ✓ $x = 10$ (6)
1.2	$3x - y + 2 = 0 \quad \text{and} \quad y = -x^2 + 2x + 8$ $\therefore y = 3x + 2 \quad \text{OR}$ $3x + 2 = -x^2 + 2x + 8 \quad 3x - (-x^2 + 2x + 8) + 2 = 0$ $x^2 + x - 6 = 0$ $(x+3)(x-2) = 0$ $x = -3 \text{ or } x = 2$ $y = 3(-3) + 2 \quad \text{or} \quad y = 3(2) + 2$ $= -7 \quad \text{or} \quad y = 8$	✓ $y = 3x + 2$ ✓ substitution/vervanging ✓ std form/stand vorm ✓ factors/faktore ✓ x -values/ x -waardes ✓ y -values/ y -waardes (6)
1.3	$3x^2 + (k+2)x = 1 - k$ $3x^2 + (k+2)x - 1 + k = 0$ $\Delta = b^2 - 4ac$ $= (k+2)^2 - 4(3)(-1+k)$ $= k^2 + 4k + 4 + 12 - 12k$ $= k^2 - 8k + 16$ $= (k-4)^2$ $\therefore b^2 - 4ac \text{ is a perfect square.}$ <p>Roots are real and rational.</p>	✓ $\Delta = b^2 - 4ac$ ✓ substitution /vervanging ✓ $k^2 - 8k + 16$ ✓ $(k-4)^2$ (4) [30]



QUESTION/VRAAG 2

2.1.1	$\begin{aligned} & \frac{5^a \cdot 5^{-2} \cdot 2^a \cdot 2^2}{10^a - 10^a \cdot 10^{-1} \cdot 2} \\ &= \frac{(5 \cdot 2)^a \cdot 5^{-2} \cdot 2^2}{10^a \left[1 - \frac{2}{10} \right]} \\ &= \frac{10^a \cdot \frac{4}{25}}{10^a \cdot \frac{8}{10}} \\ &= \frac{4}{25} \times \frac{10}{8} \\ &= \frac{1}{5} \end{aligned}$	<ul style="list-style-type: none"> ✓ writing as separate bases/ skryf as priembasisse ✓ multiplication of bases with same exponents/vermenigv. van basisse met dies. eksp. ✓ common factor in the denominator/gemene faktor in die noemer ✓ simplification/vereenv. ✓ answer/antw.
2.1.2	$\begin{aligned} & \frac{\sqrt{27m^6} - \sqrt{48m^6}}{\sqrt{12m^6}} \\ &= \frac{3\sqrt{3}m^3 - 4\sqrt{3}m^3}{2\sqrt{3m^6}} \\ &= \frac{\sqrt{3m^6}(3-4)}{2\sqrt{3}m^3} \\ &= \frac{-\sqrt{3}m^3}{2\sqrt{3}m^3} \\ &= -\frac{1}{2} \end{aligned}$ <p>OR/OF</p> $\begin{aligned} & \frac{3\sqrt{3m^6} - 4\sqrt{3m^6}}{2\sqrt{3m^6}} \\ &= \frac{\sqrt{3m^6}(3-4)}{2\sqrt{3m^6}} \\ &= \frac{3-4}{2} \\ &= -\frac{1}{2} \end{aligned}$	<ul style="list-style-type: none"> ✓ simplification of all surds/vereenv. van alle wortelvorme ✓ simplification numerator/vereenv. van teller ✓ answer/antw. <p>(3)</p> <ul style="list-style-type: none"> ✓ simplification of all surds/ vereenv. van alle wortelvorme ✓ simplification of numerator/ vereenv. van teller ✓ answer/antw. <p>(3)</p>

2.2	$\begin{aligned} \text{LHS} &= \frac{4\sqrt{2} - 8(1 + \sqrt{2})}{2\sqrt{2}(1 + \sqrt{2})} \\ &= \frac{-4\sqrt{2} - 8}{2\sqrt{2}(1 + \sqrt{2})} \\ &= \frac{-4(\sqrt{2} + 2)}{2(\sqrt{2} + 2)} \\ &= -2 \\ &= \text{RHS} \end{aligned}$ <p>OR/OF</p> $\begin{aligned} \text{LHS} &= \frac{2}{1 + \sqrt{2}} \times \frac{1 - \sqrt{2}}{1 - \sqrt{2}} - \frac{8 \times \sqrt{8}}{\sqrt{8} \times \sqrt{8}} \\ &= \frac{2 - 2\sqrt{2}}{1 - 2} - \sqrt{8} \\ &= -2 + 2\sqrt{2} - 2\sqrt{2} \\ &= -2 \\ &= \text{RHS} \end{aligned}$	<ul style="list-style-type: none"> ✓ LCD/KGV ✓ numerator/teller ✓ simplification/ vereenv. ✓ common factor/gemene faktor <p>(4)</p>
-----	--	---



Dm


QUESTION/VRAAG 3

3.1	<p>$x - 23 = 4$ $x = 27$</p>	✓ answer/antw. (1)
3.2	$2a = 4$ $a = 2$ $3a + b = 3$ $6 + b = 3$ $b = -3$ $a + b + c = -9$ $2 - 3 + c = -9$ $c = -8$ $T_n = 2n^2 - 3n - 8$	$\checkmark a = 2$ $\checkmark b = -3$ $\checkmark c = -8$ $\checkmark T_n = 2n^2 - 3n - 8$ (4)
3.3	$T_n = 2n^2 - 3n - 8 + 3$ $= 2n^2 - 3n - 5$	CA from 3.2 \checkmark answer/antw. (1)
3.4	$T_n = 400$ $2n^2 - 3n - 5 = 400$ $2n^2 - 3n - 405 = 0$ $(n-15)(2n+27) = 0$ $n = 15 \text{ or } n \neq \frac{-27}{2}$	CA from 3.3 \checkmark equating/verg. \checkmark std form/stand vorm \checkmark factorisation/fakt. $\checkmark n = 15$ (4)
	OR	
	$2n^2 - 3n - 8 + 3 = 400$ $2n^2 - 3n - 8 = 397$ $2n^2 - 3n - 405 = 0$ $(n-5)(2n+27) = 0$ $n = 15 \text{ or } n \neq \frac{-27}{2}$	\checkmark equating/verg. \checkmark std form/stand vorm \checkmark factorisation/fakt. \checkmark only/ slegs $n = 15$ (4) [10]

QUESTION/VRAAG 4

4.1.1	$T_4 = 6$	✓ answer/antw. (1)
4.1.2	$\begin{aligned} T_n &= a + (n-1)d \\ &= 18 + (n-1)(-4) \\ &= -4n + 22 \end{aligned}$	Answer only, full marks/ slegs antwoord volpunte ✓ substitution/verv. ✓ answer/antw. (2)
4.1.3	$\begin{aligned} T_n &= 22 - 4n \\ -70 &= 22 - 4n \\ -92 &= -4n \\ n &= 23 \end{aligned}$	✓ substitution/verv. ✓ answer/antw. (2)
4.1.4	$\begin{aligned} Q_{510} - Q_{509} &= T_{509} \text{ of the linear sequence} \\ &= 22 - 4 \times 509 \\ &= -2014 \end{aligned}$	✓ making association/ass. ✓ answer/antw. (2)
4.2.1	$\begin{aligned} 2a &= 2 \\ \therefore a &= 1 \\ \therefore a &> 0 \\ \therefore \text{this pattern has a minimum value/hierdie patroon het 'n minimum waarde} \\ \text{The shape of the graph will be concave up / die vorm van die grafiek is konkaaf na bo} \end{aligned}$	 ✓ value/wrde of a ✓ $a > 0$ ✓ minimum value/wrde (3)
4.2.2	$\begin{aligned} T_5 &= 29 \\ \therefore 1(5)^2 + 5b + c &= 29 \\ ie 5b + c &= 4 \dots (1) \\ \text{and } T_{17} &= 29 \\ \therefore 1(17)^2 + 17b + c &= 29 \\ ie 17b + c &= -260 \dots (2) \\ \text{solve the equations simultaneously} \\ -12b &= 264 \\ \therefore b &= -22 \\ \text{substitute in (1)} \\ ie 5(-22) + c &= 4 \\ -110 + c &= 4 \\ \therefore c &= 114 \\ \therefore T_n &= n^2 - 22n + 114 \end{aligned}$	✓✓ equations/verg. (1) & (2) ✓ value of/waarde van b ✓ value of/waarde van c ✓ answer/antwoord. (5)


OR/OF



$$\begin{aligned}
 T_n &= 1(n+p)^2 + q \\
 \text{A.O.S} &= \frac{5+17}{2} \\
 p &= 11 \\
 \therefore T_n &= 1(n-11)^2 + q \\
 29 &= 1(17-11)^2 + q \\
 \therefore q &= -7 \\
 \therefore T_n &= (n-11)^2 - 7 \\
 T_n &= n^2 - 22n + 114
 \end{aligned}$$

- ✓ axis of symmetry/simm. as
- ✓ value of/waarde van p
- ✓ substitution/verv. (17 ; 29) or/of (5 ; 29)
- ✓ value of/waarde van q
- ✓ answer/antw.

[15]



Dm

QUESTION 5

5.1

$$\begin{aligned}x &= -\frac{b}{2a} \\&= -\frac{1}{2(-2)} \\&= \frac{1}{4} \\&\therefore y = -2\left(\frac{1}{4}\right)^2 + \left(\frac{1}{4}\right) + 6\end{aligned}$$

$$y = \frac{49}{8}$$

OR

$$f(x) = -2x^2 + x + 6$$

$$-2x^2 + x + 6 = 0$$

$$(2x+3)(x-2) = 0$$

$$x = \frac{-3}{2} \text{ or}$$

$$x = 2$$

x-value of the axis of symmetry

$$x = -\frac{\frac{3}{2} + 2}{2} = \left(\frac{1}{4}\right)$$

$$f\left(\frac{1}{4}\right) = -2\left(\frac{1}{4}\right)^2 + \frac{1}{4} + 6$$

$$= \frac{49}{8}$$

OR

$$f(x) = -2\left(x^2 - \frac{x}{2}\right) + 6$$

$$= -2\left(x^2 - \frac{x}{2} + \left(-\frac{1}{4}\right)^2\right) + 6 - 2(-2)\left(-\frac{1}{4}\right)^2$$

$$= -2\left(x - \frac{1}{4}\right)^2 + \frac{49}{9}$$

$$TP = \left(\frac{1}{4}; \frac{49}{8}\right)$$

✓ substitution/verv.

✓ *x*-value/waarde

✓ substitution/verv.

✓ *y*-value/waarde

(4)


5.2

$$y = -2(0)^2 + 0 + 6$$


$$\therefore y \text{ intercept}(0; 6)$$

✓ *y*-value/waarde

(1)




DM


5.3	<p>x intercepts $0 = -2x^2 + x + 6$ $0 = 2x^2 - x - 6$ $0 = (2x + 3)(x - 2)$ $\therefore x = 2 \text{ or } x = -\frac{3}{2}$ $(2;0) \text{ and } \left(-\frac{3}{2};0\right)$</p>	<p>✓ $y = 0$ ✓ factorisation/faktorisering. ✓ ✓ x-values/waardes</p>
5.4		✓ shape/vorm ✓ x -intercepts/x-afsnitte ✓ turning point/draaipunt
5.5	$k = \frac{49}{8}$	✓✓ answer/antwoorde. (2)
5.6	New/Nuwe turning point/drpn.t $\left(\frac{9}{4}; \frac{57}{8}\right)$ Equation/verg. of h $y = -2\left(x - \frac{9}{4}\right)^2 + \frac{57}{8}$	✓✓ turning points/draaipunt ✓ equation/verg. OR/OF ✓✓✓ answer only [17]

QUESTION/VRAAG 6

6.1	$x = -3$ and $y = -1$	✓ $x = -3$ ✓ $y = -1$ (2)
6.2	$x \in \mathbb{R}; x \neq -3$ OR $x \in (-\infty; -3) \cup (-3; \infty)$	✓✓ answer/antwoord. (2)




6.3.1	<p>At B, $x = 0$</p> $\therefore y = \frac{1}{0+3} - 1$ $y = -\frac{2}{3}$ $\therefore OB = \frac{2}{3} \text{ units}$	<ul style="list-style-type: none"> ✓ substituton/vervanging. ✓ answer/antwoord.
6.3.2	<p>At A, $y = 0$</p> $0 = \frac{1}{x+3} - 1$ $1 = \frac{1}{x+3}$ $x+3 = 1$ $x = -2$ $\therefore OA = 2 \text{ units/ eenhede}$	<ul style="list-style-type: none"> ✓ substitution/vervanging. ✓ simplification/vereenv. ✓ answer/antwoorde.
6.4	$\frac{1}{x+3} - 1 = \frac{1}{2}x$ $2 - 2(x+3) = x(x+3)$ $x^2 + 3x - 2 + 2x + 6 = 0$ $x^2 + 5x + 4 = 0$ $(x+4)(x+1) = 0$ $x = -4 \text{ or } of \text{ } x = -1$ <p>when / wanneer $x = -1; y = -\frac{1}{2}$</p> <p>when / wanneer $x = -4; y = -2$</p> $\therefore C(-1; -\frac{1}{2}) \text{ and } D(-4; -2)$	<ul style="list-style-type: none"> ✓ equating the two equations/ verg. van 2 vergelykings ✓ standard form/std vorm ✓ factors/faktore. ✓ x-values/waardes ✓ co-ordinates/koördinate C ✓ co-ordinates/ koördinate D



6.5	$\frac{1}{x+3} \geq \frac{x+2}{2}$ $\frac{1}{x+3} \geq \frac{x}{2} + 1$ $\frac{1}{x+3} - 1 \geq \frac{x}{2}$ $\therefore f(x) \geq g(x)$ $\therefore x \leq -4 \text{ or } -3 < x \leq -1$	Answer only, full marks/ slegs antwoord volpunte	<ul style="list-style-type: none"> ✓ simplification/ vereenvoudig ✓ $f(x) \geq g(x)$ ✓ $x \leq -4$ ✓ $-3 < x \leq -1$
			(4) [19]

QUESTION/VRAAG 7


7.1	$q = 2$ $f(x) = 2 \cdot b^{x+1} + 2$ $20 = 2 \cdot b^{1+1} + 2$ $18 = 2 \cdot b^2$ $9 = b^2$ $b = 3$ $f(x) = 2 \cdot 3^{x+1} + 2$	<ul style="list-style-type: none"> ✓ substitution of / vervanging van $q = 2$ ✓ substitution of / vervanging van $(1; 20)$ ✓ $b^2 = 9$ 	(3)
7.2	$y = 2 \cdot 3^{-1+1} + 2$ $y = 2 \cdot 1 + 2$ $y = 4$	<ul style="list-style-type: none"> ✓ answer/antwoord. 	(1)
7.3	$m = \frac{y_2 - y_1}{x_2 - x_1}$ $= \frac{20 - 4}{1 - (-1)}$ $= 8$	<ul style="list-style-type: none"> ✓ substitution/vervanging. ✓ answer/antwoord. 	(2)
7.4	$h(x) = -2 \cdot 3^{x+1} + 2$ <p>OR/OF</p> <p>Reflected about the x-axis/ refleksie om die x-as</p> $= -2 \cdot 3^{x+1} - 2$ <p>\therefore Reflected about the asymptote</p> $h(x) = -2 \cdot 3^{x+1} - 2 + 4$ $= -2 \cdot 3^{x+1} + 2$	<ul style="list-style-type: none"> ✓ ✓ answer/antwoord. <p>OR/OF</p> <ul style="list-style-type: none"> ✓ reflection about x-axis/ refleksie om die x-as 	(2)
7.5	$y < 2$	<ul style="list-style-type: none"> ✓ answer/antwoord. 	(1) [9]



DM

QUESTION/VRAAG 8

8.1	$A = P(1-i)^n$ $= R 25\ 000 (1-0,09)^4$ $= R 17\ 143,74$	✓ $A = P(1-i)^n$ ✓ substitution/verv. ✓ answer/antw. (3)
8.2	$1 + i_{\text{eff}} = \left(1 + \frac{i_{\text{nom}}}{m}\right)^m$ $1 + i_{\text{eff}} = \left(1 + \frac{0,1235}{12}\right)^{12}$ $i_{\text{eff}} = \left(1 + \frac{0,1235}{12}\right)^{12} - 1$ $\therefore \text{Rate} = 0,13073 \times 100$ $= 13,07\%$ <p>The effective interest rate/Die effektiewe rentekoers is 13,07%</p>	✓ formula/formule. ✓ substitution//vervanging. ✓ simplification/vereenv. ✓ answer/antwoord. (4)
8.3	$A = P(1+i)^n$ $R 221\ 292,32 = R 145\ 000 \left(1 + \frac{r}{100}\right)^6$ $\sqrt[6]{\frac{R 221\ 292,32}{145\ 000}} = 1 + \frac{r}{100}$ $\frac{r}{100} = 0,07300000324$ $r = 7,3\%$	✓ correct substitution into correct formula/ korrekte vervanging in korrekte formule ✓ $n = 6$ ✓ $\sqrt[6]{\frac{R 221\ 292,32}{145\ 000}} = 1 + \frac{r}{100}$ ✓ answer/antw. (4)
8.4	$A = 15\ 000 \left(1 + \frac{0,096}{4}\right)^{12} - 5\ 000 \left(1 + \frac{0,096}{4}\right)^{10} + 3\ 500 \left(1 + \frac{0,096}{4}\right)^4$ $= R 17\ 448,46$	✓ $\frac{0,096}{4}$ ✓ $15\ 000 \left(1 + \frac{0,096}{4}\right)^{12}$ ✓ $- 5\ 000 \left(1 + \frac{0,096}{4}\right)^{10}$ ✓ $3\ 500 \left(1 + \frac{0,096}{4}\right)^4$ ✓ answer/antw. (5)



OR

T_0 to $T_{\frac{1}{2}}$,

$$A = 15000 \left(1 + \frac{0.069}{4}\right)^{4 \times \frac{1}{2}}$$

$$= R15728,64$$

$$\checkmark \frac{0,096}{4}$$

$$\mathbf{A} \text{ at } T_{\frac{1}{2}} = R15728,64 - R5000$$

$$= R10728,64$$

$$\checkmark R10728,64$$

$T_{\frac{1}{2}}$ to T_2 ,

$$A = R10728,64 \left(1 + \frac{0.096}{4}\right)^{\frac{3}{2} \times 4}$$

$$= R12369,28$$

$$\checkmark R12369,28$$

$$\mathbf{A} \text{ at } T_2 = R12369,28 + R3500$$

$$= R15869,28$$

$$\checkmark R15869,28$$


$$T_2 \text{ to } T_3 = R15869,28 \left(1 + \frac{0.096}{4}\right)^{4 \times 1}$$

$$= R17448,46$$

$$\checkmark R17448,46$$

(5)

[16]



DM

QUESTION/VRAAG 9


9.1	Given/Gegee: $P(A) = 0,2$ $P(B) = 0,5$ $P(A \text{ or } B) = 0,6$	
9.1.1	$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$ $0,6 = 0,2 + 0,5 - P(A \text{ and } B)$ $P(A \text{ and } B) = 0,1$	$\checkmark 0,6 = 0,2 + 0,5 - P(A \text{ and } B)$ $\checkmark P(A \text{ and } B) = 0,1$ (2)
9.1.2	$P(A \text{ and } B) = 0,1$ $P(A) \times P(B) = 0,2 \times 0,5$ $= 0,1$ $\therefore P(A \text{ and } B) = P(A) \times P(B)$ $\therefore A \text{ and } B \text{ are independent/ } A \text{ en } B \text{ is onafhanklik}$	$\checkmark P(A) \times P(B) = 0,1$ $\checkmark P(A \text{ and } B) = P(A) \times P(B)$ $\checkmark \text{conclusion/gevolgtrekking}$ (3)
9.2.1	$a = 15$ $b = 1$ $c = 38$ $d = 3$ $e = 37$	<div style="border: 1px solid black; padding: 5px;">$\checkmark a = 15$ $\checkmark b = 1$ $\checkmark c = 38$ $\checkmark d = 3$ $\checkmark e = 37$</div> (5)
9.2.2	$P(\text{one learner plays netball or volleyball}) = \frac{25}{100} = \frac{1}{4}$	$\checkmark 25$ $\checkmark \text{answer/antwoord}$ (2)
9.3.1	<p style="text-align: center;"> $C \quad 0,5 \quad (C ; C)$ $C \quad 0,4 \quad (C ; W)$ $W \quad 0,5 \quad (W ; C)$ $W \quad 0,3 \quad (W ; W)$ </p>	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> \checkmark branch at first level with probabilities/ eerste vertakking met waarskynlikhede </div> <div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> \checkmark branches at second level with probabilities/ tweede vertakkings met waarskynlikhede </div> <div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> \checkmark outcomes/uitkomste </div> (3)

9.3.2 $P(\text{second answer correct}) = P(C \text{ and } C) + P(W \text{ and } C)$
 $= (0,4 \times 0,5) + (0,6 \times 0,3)$
 $= 0,38$

- ✓ addition of probabilities/
som van waarskynlikhede
- ✓ substitution/vervanging
- ✓ answer/antwoord.

(3)
[18]**QUESTION/VRAAG 10**

10

Let one of the equal sides = x / Laat een van die sye = x the other side = $14 - 2x$ / die ander sy = $14 - 2x$

$$\begin{aligned} \text{Area} &= (14 - 2x)x \\ &= -2x^2 + 14x \\ x &= \frac{-14}{2(-2)} \\ &= \frac{7}{2} \text{ m} \\ y &= 7 \text{ m} \end{aligned}$$

- ✓ area formula/oppervl.for.

$$x = \frac{-14}{2(-2)}$$

- ✓ answer for/
antwoord van x
- ✓ answer for / antwoord van y

(4)

OR/OF \therefore the other side = $14 - 2x$ / die ander sy = $14 - 2x$


$$\begin{aligned} \therefore \text{Area} &= (14 - 2x)x \\ &= -2(x^2 - 7x) \\ &= -2\left(x^2 - 7x + \frac{49}{4} - \frac{49}{4}\right) \\ &= -2\left[\left(x - \frac{7}{2}\right)^2 - \frac{49}{4}\right] \\ &= -2\left(x - \frac{7}{2}\right)^2 + \frac{49}{2} \\ \therefore \text{when } x &= \frac{7}{2} \text{ metres it will have a maximum area} \\ \therefore \text{the other side} &= 14 - 2\left(\frac{7}{2}\right) \\ &= 7 \text{ metres} \end{aligned}$$


- ✓ area formula/oppervl.for.

- ✓ completing the square/
voltooiing van vierkant

- ✓ answer for/antwoord van x
- ✓ answer for / antwoord van y

(4)

OR/OF



Let the length be y

Width be $\frac{14-y}{2}$

$$\text{Area} = y \left(7 - \frac{1}{2}y \right)$$

$$= \frac{-1}{2}y^2 + 7y$$

$$y = \frac{-7}{2\left(\frac{-1}{2}\right)}$$

$$= 7m$$

$$\text{width} = 3,5m$$

✓ area formula/oppervl.for.

$$\checkmark y = \frac{-7}{2\left(\frac{-1}{2}\right)}$$

✓ answer for y
✓ answer for width/
antwoord van breedte

(4)

OR

$$\text{Area} = y \left(7 - \frac{1}{2}y \right)$$

$$= \frac{-1}{2}y^2 + 7y$$

$$= \frac{-1}{2}(y^2 - 14y)$$

$$= \frac{-1}{2}(y-7)^2 + \frac{49}{2}$$

$$\text{length} = 7m$$


$$\text{width} = 3,5m$$

✓ area formula/oppervl.for.

✓ completing the square/
voltooiing van die vierkant

✓ answer for /antwoord van
 y
✓ answer for width/
antwoord van breedte

(4)
[4]



TOTAL/TOTAAL:

150

Dm