



NATIONAL SENIOR CERTIFICATE EXAMINATION
NOVEMBER 2017

MATHEMATICS: PAPER I

MARKING GUIDELINES

Time: 3 hours

150 marks

These marking guidelines are prepared for use by examiners and sub-examiners, all of whom are required to attend a standardisation meeting to ensure that the guidelines are consistently interpreted and applied in the marking of candidates' scripts.

The IEB will not enter into any discussions or correspondence about any marking guidelines. It is acknowledged that there may be different views about some matters of emphasis or detail in the guidelines. It is also recognised that, without the benefit of attendance at a standardisation meeting, there may be different interpretations of the application of the marking guidelines.


SECTION A

QUESTION 1

(a) (1) $(x-1)^2 = 2(1-x)$
 $(x-1)^2 = -2(x-1)$
 $(x-1)^2 + 2(x-1) = 0$
 $(x-1)(x-1+2) = 0$
 $(x-1)(x+1) = 0$
 $x = 1 \quad x = -1$ (4)

(2) $5^{-x} \cdot 5^{x-2} = \frac{25^{2x}}{5}$
 $5^{-x} \cdot 5^{x-2} = \frac{5^{4x}}{5^1}$
 $5^{-x+x-2} = 5^{4x-1}$
 $-2 = 4x - 1$
 $x = -\frac{1}{4}$ (4)

(b) $(x+1)^2 < 9$
 $\therefore x^2 + 2x + 1 < 9$
 $\therefore x^2 + 2x - 8 < 0$
 $\therefore (x+4)(x-2) < 0$
 Critical Values: $-4; 2$



Solution: $\{x : -4 < x < 2\}$


Alternative

$(x+1)^2 < 9$
 $\therefore -3 < x+1 < 3$
 $\therefore -4 < x < 2$ (4)

(c) $(x-2)(x+4) = 0$
 $x^2 + 2x - 8 = 0$
 $\therefore b = 2$ and $c = -8$ (3)

Alternative

$(x+1)^2 < 9$
 $x^2 + 2x - 8 < 0$
 Sketch: $y = x^2 + 2x - 8$
 x -int: $x = -4; x = 2$



Solution: $\{x : -4 < x < 2\}$

(d) (1) $x - 2 = \frac{-4}{x - 2} - 4$ let $x - 2 = y$
 $y = -\frac{4}{y} - 4$ LCD: y
 $y^2 = -4 - 4y$
 $\therefore y^2 + 4y + 4 = 0$ (2)

(2) $(y + 2)^2 = 0$
 $\therefore y = -2$
 Roots are real and equal.

Alternative


$y^2 + 4y + 4 = 0$
 $\therefore \Delta = 4^2 - 4(1)(4)$
 $\therefore \Delta = 0$
 \therefore Roots are real and equal.

Alternative

$x - 2 = \frac{-4}{x - 2} - 4$
 $\therefore (x - 2)^2 = -4 - 4(x - 2)$
 $\therefore x^2 - 4x + 4 = -4 - 4x + 8$
 $\therefore x^2 = 0$
 \therefore Roots are real and equal.

(2)
[19]

QUESTION 2




(2) $E = \{HTT, THT, TTH\}$
 $\therefore P(2 \text{ tails and } 1 \text{ head}) = \frac{3}{8}$ (2)

(b) (1) $P(A \cap B) = 0$ (1)

(2) (i) You cannot pick a R2 and a R5 coin at the same time. (1)

(ii) $P(\text{either a R5 or a R2})$
 $= P(A \text{ or } B)$
 $= P(A) + P(B) \text{ mutually exclusive}$
 $= 0,36 + 0,47$
 $= 0,83$ (3)

(c) (1)



(4)

(2) $P(\text{exactly one machine is stamping R5 coins})$
 $= 0,3 + 0,2$
 $= 0,5$
 $\therefore 50\%$ (3)

[17]

QUESTION 3

(a) $480\ 163 \div 0,502 = R956\ 500$ (2)

(b) $R956\ 500 \times 5\% = R47\ 825$ (2)

(c) Cost of machinery including import charges = $R956\ 500 + R47\ 825$
 = $R1\ 004\ 325$

$$A = P (1 + i)^n$$

$$1\ 004\ 325 = 225\ 450 \left(1 + \frac{9,5}{100}\right)^n$$

$$\frac{1\ 004\ 325}{225\ 450} = \left(\frac{219}{200}\right)^n$$

$$\log_{\left(\frac{219}{200}\right)} \left(\frac{1\ 004\ 325}{225\ 450}\right) = n$$

$$n = 16,46171594$$

$$\therefore n \approx 16,46$$

$$\therefore \text{approx. 17 years} \quad (4)$$

(d) (1) Loan required: $R1\ 004\ 325 - R225\ 450 = R778\ 875$

$$P = x \left[\frac{1 - (1 + i)^{-n}}{i} \right]$$

$$778\ 875 = x \left[\frac{1 - \left(1 + \frac{12}{1200}\right)^{(-4 \times 12)}}{\frac{12}{1200}} \right]$$

$$x = R20\ 510,76607$$

$$\therefore x = R20\ 510,77 \quad (4)$$

(2) Outstanding Balance = $A - F$

$$A = 778\,875 \left(1 + \frac{12}{1200} \right)^{24}$$

$$A = 988\,964,5744$$

$$A \approx 988\,964,57$$

$$F = 20\,510,76607 \left[\frac{\left(1 + \frac{12}{1200} \right)^{24} - 1}{\frac{12}{1200}} \right]$$

$$F = 553\,246,4277$$

$$F \approx 553\,246,43$$

$$\text{Outstanding balance} = 988\,964,5744 - 553\,246,4277$$

$$= R435\,718,1467 \approx R435\,718,15$$

NB: If A and F are rounded to the nearest cent, consider

$$\text{Outstanding balance} = 988\,964,57 - 553\,246,43$$

$$= R435\,718,14$$

Alternative


$$\text{Outstanding balance} = 20\,510,76607 \left[\frac{1 - \left(1 + \frac{12}{1200} \right)^{-24}}{\frac{12}{1200}} \right]$$

$$= R435\,718,1466$$

$$\approx R435\,718,15$$

(3)
[15]

QUESTION 4



Constant second difference (1)

(2) $T_n = an^2 + bn + c$
 $T_1 = a + b + c = 1$
 $T_2 = 4a + 2b + c = 3$
 $T_3 = 9a + 3b + c = 6$

$\therefore 3a + b = 2$ and $5a + b = 3$
 Substitute $b = 2 - 3a$ into $5a + b = 3$
 $\therefore 5a + (2 - 3a) = 3$
 $2a = 1$
 $a = \frac{1}{2}$

$b = \frac{1}{2}$ and $c = 0$

$\therefore T_n = \frac{1}{2}n^2 + \frac{1}{2}n$ (6)

(b) $T_3 = 52$ cm
 $T_7 = 78$ cm
 $T_3 = a + 2d = 52$
 $T_7 = a + 6d = 78$

$4d = 26 \therefore d = 6\frac{1}{2}$

$\therefore a = 39$ cm

$T_{43} = 39 + 42\left(6\frac{1}{2}\right)$

$T_{43} = 312$ cm

(5)
[12]

QUESTION 5

(a) (1) $f(x) = x^2 - 6x + 9$

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^2 - 6(x+h) + 9 - (x^2 - 6x + 9)}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - 6x - 6h + 9 - x^2 + 6x - 9}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{2xh + h^2 - 6h}{h}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{h(2x + h - 6)}{h}$$

$$f'(x) = 2x - 6$$

(5)

(2) $f'(-3) = 2(-3) - 6 = -12$

(2)

(b) $y = \pi x^{-1} + 3x^{\frac{1}{3}}$

$$\frac{dy}{dx} = -\pi x^{-2} + x^{-\frac{2}{3}}$$

(4)

[11]**74 marks**

SECTION B

QUESTION 6

- (a) (1) Domain = $x \in \mathbb{R} ; x \neq 3$ (1)
 (2) Range = $y \in \mathbb{R} ; y \neq -3$ (1)
 (3) (i) 5 units (1)
 (ii) 5 units (1)

- (b) (1) $y = a \cdot b^x$ substitute $\left(0; \frac{1}{4}\right)$


$$\frac{1}{4} = a \cdot b^0$$

$$a = \frac{1}{4}$$
 $y = \frac{1}{4} b^x$ substitute $\left(2; \frac{9}{4}\right)$

$$\frac{9}{4} = \frac{1}{4} b^2$$

$$b^2 = 9$$

$$\therefore b = \pm 3 \text{ but } b > 0 \therefore b = 3$$
 (4)



(3) Range = $\left[\frac{1}{4}; \infty\right)$ (1)

(4) $f(x) = \frac{1}{4} \cdot 3^x$
 For f^{-1} : $x = \frac{1}{4} \cdot 3^y ; y \geq 0$
 $4x = 3^y$
 $y = \log_3(4x)$ for $x \geq \frac{1}{4}$ (3)

(5) See graph in Question 6 (b) (2) above. (3)

[18]

QUESTION 7

$$\begin{aligned}
 \text{(a)} \quad f(x) &= x^2 + 6x + (3)^2 + 5 - (3)^2 \\
 f(x) &= (x+3)^2 - 4 \\
 \therefore \text{T.P.} &(-3; -4)
 \end{aligned}
 \tag{3}$$

$$\begin{aligned}
 \text{(b)} \quad (1) \quad x^2 + 6x + 5 &= -x - 5 \\
 x^2 + 7x + 10 &= 0 \\
 x &= -2 \text{ or } x = -5 \\
 A(-5; 0) \text{ and } B(-2; -3)
 \end{aligned}
 \tag{4}$$

$$(2) \quad \text{Horizontal shift: } \therefore -5 < t < -2 \tag{3}$$

$$\begin{aligned}
 \text{(c)} \quad (1) \quad \text{Length MN} &= (-x - 5) - (x^2 + 6x + 5) \\
 \text{Length MN} &= -x - 5 - x^2 - 6x - 5 \\
 \text{Length MN} &= -x^2 - 7x - 10
 \end{aligned}$$

For max. length: Let $D_x = 0$

$$-2x - 7 = 0$$

$$x = -\frac{7}{2}$$

$$\therefore \text{Max. length MN} = -\left(-\frac{7}{2}\right)^2 - 7\left(-\frac{7}{2}\right) - 10$$

$$\therefore \text{Max. length MN} = \frac{9}{4} \text{ units} \quad \text{i.e. } 2,25 \text{ units} \tag{6}$$

$$(2) \quad \text{Vertical shift: } \therefore k > \frac{9}{4} \tag{1}$$

[17]

QUESTION 8

(a) (1) $\frac{3}{2}; -\frac{9}{2}; \frac{27}{2}; \dots$
 $\therefore r = -3$ and series is geometric
 however, series is not convergent since $r < -1$.
 $\therefore x \neq -\frac{3}{2}$ (3)

(2) $\frac{x-3}{x+3} = \frac{12-x}{x-3}$
 $(x-3)^2 = (12-x)(x+3)$
 $x^2 - 6x + 9 = 12x + 36 - x^2 - 3x$
 $2x^2 - 15x - 27 = 0$
 $x = 9$ or $x \neq -\frac{3}{2}$ (4)

(b) $S_4 = 7\frac{1}{2}; S_5 = 15\frac{1}{2}$ and $S_6 = 31\frac{1}{2}$
 $T_5 = S_5 - S_4$
 $T_5 = 8$

 $T_6 = S_6 - S_5$
 $T_6 = 16$

 $T_5 = ar^4 = 8$
 $T_6 = ar^5 = 16$

 $\frac{T_6}{T_5} = r = 2$
 $\therefore a = \frac{1}{2}$
 $S_n = \frac{\frac{1}{2}(2^n - 1)}{2 - 1}$
 $= 2^{n-1} - \frac{1}{2}$ (7)

[14]

QUESTION 9

(a) $f(x) = -x^3 + bx^2 + cx - 3$
 $f(1) = -(1)^3 + b(1)^2 + c(1) - 3 = 4$
 $b + c = 8$

$$f'(x) = -3x^2 + 2bx + c$$

$$f''(x) = -6x + 2b$$

$$f''\left(\frac{1}{2}\right) = -6\left(\frac{1}{2}\right) + 2b = 1$$

$$b = 2$$

$$\therefore c = 6$$

(7)

(b) For concave up: $f''(x) > 0$

$$-6x + 4 > 0$$

$$x < \frac{2}{3}$$

(3)

[10]

QUESTION 10

$$\frac{340}{x} - \frac{340}{x+2} = 3 \quad \text{LCD: } x(x+2)$$

$$340(x+2) - 340x = 3x(x+2)$$

$$3x^2 + 6x - 680 = 0$$

$$x = 14,09 \quad \text{or} \quad x \neq -16,09$$

$$\text{Therefore Time} = \frac{340}{14,09} \approx 24,13 \text{ seconds}$$

Alternative

Let original time taken be represented by y .

$$\therefore xy = 340 \quad \dots \text{ eq. 1}$$

$$(x+2)(y-3) = 340 \quad \dots \text{ eq. 2}$$

$$\text{From eq. 1} \quad y = \frac{340}{x}$$

$$\therefore (x+2) \left(\frac{340}{x} - 3 \right) = 340$$

$$\therefore 3x^2 + 6x - 680 = 0$$


$$\therefore x = 14,09 \quad \text{or} \quad x \neq -16,09$$

$$\text{Therefore Time} = \frac{340}{14,09} \approx 24,13 \text{ seconds}$$

[6]

QUESTION 11

(a) (1) When $x = -2$ and $x = 0$ (2)



(2)

(b) $y = \frac{1}{5}x^3 + \frac{3}{4}x + 3$
 $\frac{dy}{dx} = \frac{3}{15}x^2 + \frac{3}{4}$ substitute $x = 0$
 $\frac{dy}{dx} = \frac{3}{4}$

Equation of tangent: $y = \frac{3}{4}x + c$ where $c = 3$

Equation of tangent: $y = \frac{3}{4}x + 3$

For point of intersection between tangent

and line BC, substitute $x = 2$ into $y = \frac{3}{4}x + 3$

$$\therefore y = 4\frac{1}{2} \therefore \text{Pt}\left(2; 4\frac{1}{2}\right)$$

Area of Busi's region = $\frac{1}{2}\left(5 + 3\frac{1}{2}\right) \times 2$

$$= 8\frac{1}{2} \text{ units}^2$$

Area of Khanya's region = $\frac{1}{2}\left(3 + 4\frac{1}{2}\right) \times 2$

$$= 7\frac{1}{2} \text{ units}^2$$

Therefore Busi's region is larger.

(7)
[11]

76 marks

Total: 150 marks